

JEE (ADVANCED) 2023 PAPER-2

[PAPER WITH SOLUTION]



HELD ON SUNDAY 04TH JUNE 2023

CHEMISTRY


SECTION 1 (Maximum Marks : 12)

- This section contains **FOUR (04)** questions.
- Each question has **FOUR** options (A), (B), (C) and (D). **ONLY ONE** of these four options is the correct answer.
- For each question, choose the option corresponding to the correct answer.
- Answer to each question will be evaluated according to the following marking scheme:
 Full Marks : +3 If **ONLY** the correct option is chosen;
 Zero Marks : 0 If none of the options is chosen (i.e. the question is unanswered);
 Negative Marks : -1 In all other cases.


[Q.1] The correct molecular orbital diagram for F_2 molecule in the ground state is




[:C]



[:D]



[:ANS] C

[:SOLN] Energy order of M.O for F_2 is

$$\sigma(1s^2) < \sigma^*(1s^2) < \sigma(2s^2) < \sigma^*(2s^2) < \sigma(2p_z^2) < \pi(2p_x^2 = 2p_y^2) < \pi^*(2p_x^2 = 2p_y^2) < \sigma^*(2p_z^0)$$

[:Q.2] Consider the following statements related to colloids.

- (I) Lyophobic colloids are not formed by simple mixing of dispersed phase and dispersion medium.
- (II) For emulsions, both the dispersed phase and the dispersion medium are liquid.
- (III) Micelles are produced by dissolving a surfactant in any solvent at any temperature.
- (IV) Tyndall effect can be observed from a colloidal solution with dispersed phase having the same refractive index as that of the dispersion medium.

The option with the correct set of statements is

[A] (I) and (II)

[B] (II) and (III)

[C] (III) and (IV)

[D] (II) and (IV)

[ANS] [A] (I) & (II)

[SOLN] Lyophobic colloids are formed by special methods of preparation.





Emulsion : (i) Dispersed phase – Liquid

(ii) Dispersion medium - Liquid

Micelles are produced by dissolving a surfactant in any solvent at or above a particular temperature, called kraft temperature

Tyndall effect can be observed from a colloidal solution with dispersed phase having refractive index different from dispersion medium.

[Q.3] In the following reactions, **P**, **Q**, **R**, and **S** are the major products



The correct statement about **P**, **Q**, **R**, and **S** is


[A] **P** is a primary alcohol with four carbons]

[B] **Q** undergoes Kolbe's electrolysis to give an eight-carbon product.



[C] **R** has six carbons and it undergoes Cannizzaro reaction.

[D] **S** is a primary amine with six carbons.


[:ANS] B



[:SOLN]







It does not give Cannizaro reaction



It's secondary amine

[Q.4] A disaccharide X cannot be oxidised by bromine water. The acid hydrolysis of X leads to a laevorotatory solution. The disaccharide X is



[ANS] A

[SOLN] B and C having hemiacetal group

∴ B and C are oxidised by Bromine water

A is structure of sucrose.

Hydrolysis of sucrose gives dextrorotatory glucose and levorotatory fructose but overall it gives levorotatory solution

SECTION 2 (Maximum Marks : 12)

- This section contains **THREE (03)** questions.
- Each question has **FOUR** options (A), (B), (C) and (D). **ONE OR MORE THAN ONE** of these four option(s) is (are) correct answer(s).
- For each question, choose the option(s) corresponding to (all) the correct answer(s).
- Answer to each question will be evaluated according to the following marking scheme:
 Full Marks : +4 **ONLY** if (all) the correct option(s) is(are) chosen;
 Partial Marks : +3 If all the four options are correct but **ONLY** three options are chosen;
 Partial Marks : +2 If three or more options are correct but **ONLY** two options are chosen, both of which are correct;
 Partial Marks : +1 If two or more options are correct but **ONLY** one option is chosen and it is a correct option;
 Zero Marks : 0 If unanswered;
 Negative Marks : -2 In all other cases.
- For example, in a question, if (A), (B) and (D) are the **ONLY** three options corresponding to correct answers, then choosing **ONLY** (A), (B) and (D) will get +4 marks; choosing **ONLY** (A) and (B) will get +2 marks;
 choosing **ONLY** (A) and (D) will get +2marks;
 choosing **ONLY** (B) and (D) will get +2 marks;
 choosing **ONLY** (A) will get +1 mark;
 choosing **ONLY** (B) will get +1 mark;
 choosing **ONLY** (D) will get +1 mark;
 choosing no option(s) (i.e. the question is unanswered) will get 0 marks and
 choosing any other option(s) will get -2 marks.

[:Q.5] The complex(es), which can exhibit the type of isomerism shown by $[\text{Pt}(\text{NH}_3)_2\text{Br}_2]$, is/are [en = $\text{H}_2\text{NCH}_2\text{CH}_2\text{NH}_2$]

[:A] $[\text{Pt}(\text{en})(\text{SCN})_2]$

[:B] $[\text{Zn}(\text{NH}_3)_2\text{Cl}_2]$



[:C] $[\text{Pt}(\text{NH}_3)_2\text{Cl}_4]$

[:D] $[\text{Cr}(\text{en})_2(\text{H}_2\text{O})(\text{SO}_4)]^+$



[:ANS] **C, D**

[:SOLN] $[\text{Pt}(\text{NH}_3)_2(\text{Br})_2]$ shows geometrical isomerism.


$[\text{Pt}(\text{NH}_3)_2(\text{Cl})_4]$ & $[\text{Cr}(\text{en})_2(\text{H}_2\text{O})(\text{SO}_4)]^+$ also shows geometrical isomerism.




(B) Tetrahedral complex do not show G.I.



(C)



(D)



[:Q.6] Atoms of metals x, y, and z from face-centred cubic (fcc) unit cell of edge length L_x , body-centred cubic (bcc) unit cell of edge length L_y , and simple cubic unit cell of edge length L_z , respectively.

If $r_z = \frac{\sqrt{3}}{2} r_y$, $r_y = \frac{8}{\sqrt{3}} r_x$, $M_z = \frac{3}{2}$ and $M_z = 3M_x$, then the correct statement(s) is(are)

[Given: M_x , M_y , and M_z are molar masses of metals x, y, and z, respectively.

r_x , r_y , and r_z are atomic radii of metals x, y, and z respectively.

[:A] Packing efficiency of unit cell of x > Packing efficiency of unit of y > Packing efficiency of unit cell of z

[:B] $L_y > L_z$

$$[:C] \quad L_x > L_y$$

$$[:D] \quad \text{Density of } x > \text{Density of } y$$

$$[:ANS] \quad A, B, D$$

$$[:SOLN] \quad \text{FCC (for } x)$$

$$\text{edge length} = L_x$$

relation

$$4r_x = \sqrt{2} L_x$$

$$r_x = \frac{\sqrt{2} L_x}{4} \quad \dots(i)$$

BCC (for } y)

$$\text{edge length} = L_y$$

relation

$$4r_y = \sqrt{3} L_y$$

$$r_y = \frac{\sqrt{3} L_y}{4} \quad \dots(2)$$

SC (for } Z)

$$\text{edge length} = L_z$$

$$2r_z = L_z$$

$$r_z = \frac{L_z}{2} \quad \dots(3)$$

eqn (1) \div eqn (2)

$$\frac{r_x}{r_y} = \frac{\sqrt{2} L_x \times 4}{4 \times \sqrt{3} L_y}$$

$$\frac{r_x}{r_y} = \frac{\sqrt{2} L_x}{\sqrt{3} L_y}$$

$$\frac{8}{\sqrt{3}} = \frac{\sqrt{3} L_y}{\sqrt{2} L_x}$$

$$\frac{L_x}{L_y} = \frac{\sqrt{3} \times \sqrt{3}}{\sqrt{2} \times 8} = \frac{3}{\sqrt{2} \times 8}$$

$$\frac{L_x}{L_y} < 1 \quad \text{So, } L_y > L_x$$

So, 'C' wrong

eqn (3) ÷ eqn (2)

$$\frac{r_z}{r_y} = \frac{L_z \times 4}{2 \times \sqrt{3} L_y}$$

$$\frac{\sqrt{3}}{2} = \frac{4}{2\sqrt{3}} \times \frac{L_z}{L_y}$$

$$\frac{3}{4} = \frac{L_z}{L_y}$$

$$\frac{L_y}{L_z} = \frac{4}{3}$$

$$\frac{L_y}{L_z} > 1 \quad \text{So, } L_y > L_z$$

$$D_x = \frac{Z_x \times M_x}{N_A \times L_x^3}$$

$$D_y = \frac{Z_y \times M_y}{N_A \times L_y^3}$$

$$\frac{D_x}{D_y} = \frac{Z_x \times M_x \times \cancel{N_A} \times L_y^3}{\cancel{N_A} \times L_x^3 \times Z_y \times M_y}$$

$$\frac{D_x}{D_y} = \left(\frac{Z_x}{Z_y} \right) \times \left(\frac{M_x}{M_y} \right) \times \left(\frac{L_y}{L_x} \right)^3$$

$$\frac{D_x}{D_y} = \frac{4}{2} \times \frac{1}{2} \times \left(\frac{3}{\sqrt{3} \times 8} \right)^3 \left\{ \begin{array}{l} \frac{3}{2} M_y = 3 M_n \\ \frac{1}{2} = \frac{M_x}{M_y} \end{array} \right.$$

$$\frac{D_y}{D_x} = \frac{27}{2\sqrt{2} \times 512}$$

$$\frac{D_x}{D_y} = \frac{2\sqrt{2} \times 512}{27}$$

$$D_n > D_y$$

$$p.f = \frac{Z \times \frac{4}{3} \pi r^3}{a^3}$$

for FCC $4r = \sqrt{2}a$, $Z = 4$

$$\text{p.f.} = \frac{4 \times \frac{4}{3} \pi r^3}{\left(\frac{4r}{\sqrt{2}}\right)^3} = 0.74 \quad \{\text{Independent from } L_x \text{ \& } r_n\}$$

For BCC

$$4r = \sqrt{3}a \quad Z = 2$$




$$\text{p.f.} = \frac{2 \times \frac{4}{3} \pi r^3}{\left(\frac{4r}{\sqrt{3}}\right)^3} = 0.68 \quad \{\text{Independent from } L_n \text{ \& } r_x\}$$


For SC $2r = a$ $Z = 1$

$$\text{p.f.} = \frac{1 \times \frac{4}{3} \pi r^3}{(2r)^3} = 0.52 \quad \{\text{Independent from } L_x \text{ \& } r_x\}$$

So, p.f. for x > p.f. for y > p.f. for z

[:Q.7] In the following reactions, **P**, **Q**, **R**, and **S** are the major products.





The correct statement(s) about P, Q, R, and S is(are)

[A] **P** and **Q** are monomers of polymers dacron and glyptal, respectively.




[B] **P**, **Q**, and **R** are dicarboxylic acids.


[C] Compounds **Q** and **R** are the same.

[D] **R** does not undergo aldol condensation and **S** does not undergo Cannizzaro reaction.

[ANS] C,D

[SOLN]






SECTION 3 (Maximum Marks : 24)


- This section contains **SIX (06)** questions.
- The answer to each question is a **NON-NEGATIVE INTEGER**.
- For each question, enter the correct integer corresponding to the answer using the mouse and the onscreen virtual numeric keypad in the place designated to enter the answer.
- Answer to each question will be evaluated according to the following marking scheme:
Full Marks : +4 If **ONLY** the correct integer is entered;
Zero Marks : 0 In all other cases.


[:Q.8] H_2S (5 moles) reacts completely with acidified aqueous potassium permanganate solution. In this reaction, the number of moles of water produced is x , and the number of moles of electrons involved is y . The value of $(x + y)$ is _____.

[:ANS] 18



[:SOLN] $[\text{H}_2\text{S} + [\text{O}] \rightarrow \text{H}_2\text{O} + \text{S} \downarrow] \times 5$





Total no of moles of H₂O produced = 8 (x)


Total no. of moles of electron involved = 10 (y)


$$\therefore (x + y) = 18$$

[:Q.9] Among $[\text{I}_3]^+$, $[\text{SiO}_4]^{4-}$, SO_2Cl_2 , XeF_2 , SF_4 , ClF_3 , $\text{Ni}(\text{CO})_4$, XeO_2F_2 , $[\text{PtCl}_4]^{2-}$, XeF_4 , and SOCl_2 , the total number of species having sp³ hybridised central atom is _____.

[:ANS] 5

[:SOLN]










[:Q.10] Consider the following molecules: Br_3O_8 , F_2O , $\text{H}_2\text{S}_4\text{O}_6$, $\text{H}_2\text{S}_5\text{O}_6$, and C_3O_2 .
Count the number of atoms existing in their zero oxidation state in each molecule.
Their sum is _____

[:ANS] 6

[:SOLN]



[Q.11] For He^+ , a transition takes place from the orbit of radius 105.8 pm to the orbit of radius 26.45 pm. The wavelength (in nm) of the emitted photon during the transition is ____.

[Use:

Bohr radius, $a = 52.9$ pm

Rydberg constant, $R_H = 2.2 \times 10^{-18}$ J

Planck's constant, $h = 6.6 \times 10^{-34}$ J s

Speed of light, $c = 3 \times 10^8$ m s^{-1}]

[ANS] $\lambda = 30\text{nm}$

[SOLN] $\frac{52.9 \times n_2^2}{2} = 105.8$

$$n_2 = 2$$

Now

$$\frac{52.9 \times n_1^2}{2} = 26.45$$

$$n_1 = 1$$

$$\Delta E = 2.2 \times 10^{-18} Z^2 \left(\frac{1}{n_1^2} - \frac{1}{n_2^2} \right) = \frac{hc}{\lambda}$$

$$2.2 \times 10^{-18} \times 4 \times \frac{3}{4} = \frac{hc}{\lambda}$$

$$2.2 \times 10^{-18} \times 3 = \frac{6.6 \times 10^{-34} \times 3 \times 10^8}{\lambda}$$

$$\lambda = 30 \times 10^{-9} \text{ metre}$$

$$\lambda = 30\text{nm}$$

[Q.12] 50 mL of 0.2 molal urea solution (density = 1.012 g mL^{-1} at 300 K) is mixed with 250 mL of solution containing 0.06 g of urea. Both the solutions were prepared in the same solvent. The osmotic pressure (in Torr) of the resulting solution at 300 K is ____.

[Use: Molar mass of urea = 60 g mol^{-1} ; gas constant, $R = 62$ L Torr K^{-1} mol^{-1} ;

Assume, $\Delta_{\text{mix}} H = 0$, $\Delta_{\text{mix}} V = 0$]

[ANS] 682 torr

[SOLN] 0.2 mole solute in 1000 g solvent

solute = 12 g

solution = 1012 g

$$\text{volume of solution} = \frac{1012}{1.012} \text{ g}$$

1000 g

$$M_1 = 0.2 \times \frac{1000}{1000} = 0.2$$

$$M_2 = \frac{0.06}{60} \times \frac{1000}{250} = \frac{1}{250}$$

$$M_1 V_1 + M_2 V_2 = M_R V_R$$

$$0.2 \times 50 + \frac{1}{250} \times 250 = M_R \times 300$$

$$M_R = \frac{11}{300}$$

$$\pi = CRT$$

$$= \frac{11}{300} \times 62 \times 300$$


$$= 682 \text{ torr}$$


[Q.13] The reaction of 4-methyloct-1-ene (**P**, 2.52 g) with HBr in the presence of $(\text{C}_6\text{H}_5\text{CO})_2\text{O}_2$ gives two isomeric bromides in a 9 : 1 ratio, with a combined yield of 50%. Of these, the entire amount of the primary alkyl bromide was reacted with an appropriate amount of diethylamine followed by treatment with aq. K_2CO_3 to give a non-ionic product **S** in 100% yield.

The mass (in mg) of **S** obtained is ____.

[Use molar mass (in g mol^{-1}): H = 1, C = 12, N = 14, Br = 80]

[ANS] 1791






SECTION 4 (Maximum Marks : 12)

- This section contains **TWO (02)** paragraphs.
- Based on each paragraph, there are **TWO (02)** questions.
- The answer to each question is a **NUMERICAL VALUE**.
- For each question, enter the correct numerical value of the answer using the mouse and the onscreen virtual numeric keypad in the place designated to enter the answer.
- If the numerical value has more than two decimal places, **truncate/round-off** the value to **TWO** decimal places.
- Answer to each question will be evaluated according to the following marking scheme:
 Full Marks : +3 If **ONLY** the correct numerical value is entered in the designated place;
 Zero Marks : 0 In all other cases.

Passage-1

The entropy versus temperature plot for phases α and β at 1 bar pressure is given.

S_T and S_0 are entropies of the phases at temperatures T and 0 K, respectively



The transition temperature for α to β phase change is 600 K and $C_{p,\beta} - C_{p,\alpha} = 1 \text{ J mol}^{-1} \text{ K}^{-1}$. Assume $(C_{p,\beta} - C_{p,\alpha})$ is independent of temperature in the range of 200 to 700 K. $C_{p,\alpha}$ and $C_{p,\beta}$ are heat capacities of α and β phases, respectively.

[Q.14] The value of entropy change, $S_\beta - S_\alpha$ (in $\text{J mol}^{-1} \text{ K}^{-1}$), at 300 K is _____.

[Use : $\ln 2 = 0.69$

Given : $S_\beta - S_\alpha = 0$ at 0 K]

[ANS] 0.31

[SOLN] $d(\Delta S) = \frac{\Delta C_p dT}{T}$

$$\int_{T_1}^{T_2} d\Delta S = \Delta C_p \int_{T_1}^{T_2} \frac{dT}{T}$$

$$\Delta S_{T_2} - \Delta S_{T_1} = \Delta C_p \ln \frac{T_2}{T_1}$$

$$(S_{600} - S_0) - (S_{300} - S_0) = \Delta C_p \ln \frac{T_2}{T_1}$$

$$\Delta S_{600} - \Delta S_{300} = \Delta C_p \ln \frac{600}{300}$$

$$1 - \Delta S_{300} = 1 \times \ln 2 = 0.69$$


$$\Delta S_{300} = 1 - 0.69$$

$$\Delta S_{300} = 0.31$$

Passage-1

The entropy versus temperature plot for phases α and β at 1 bar pressure is given.

S_T and S_0 are entropies of the phases at temperatures T and 0 K, respectively



The transition temperature for α to β phase change is 600 K and $C_{p,\beta} - C_{p,\alpha} = 1 \text{ J mol}^{-1} \text{ K}^{-1}$. Assume $(C_{p,\beta} - C_{p,\alpha})$ is independent of temperature in the range of 200 to 700 K. $C_{p,\alpha}$ and $C_{p,\beta}$ are heat capacities of α and β phases, respectively.

[:Q.15] The value of enthalpy change, $H_\beta - H_\alpha$ (in J mol^{-1}), at 300 K is _____.

[:ANS] 300

[:SOLN]
$$\int_{T_1}^{T_2} d(\Delta H) = \Delta C_p \int_{T_1}^{T_2} dT$$

$$\Delta H_{T_2} - \Delta H_{T_1} = \Delta C_p (T_2 - T_1)$$

$$\Delta H_{600} - \Delta H_{300} = \Delta C_p (600 - 300) \quad \dots(1)$$

$$\Delta S_{600} = \frac{\Delta H_{600}}{600} \Rightarrow \Delta H_{600} = 600 \Delta S_{600}$$

$$= 600$$

From eqn (i)

$$600 - \Delta H_{300} = 1 \times 300 \Rightarrow \Delta H_{300} = 300$$

Passage-2

A trinitro compound, 1,3,5-tris-(4-nitrophenyl)benzene, on complete reaction with an excess of Sn/HCl gives a major product, which on treatment with an excess of NaNO₂/HCl at 0 °C provides P as the product. P, upon treatment with excess of H₂O at room temperature, gives the product Q. Bromination of Q in aqueous medium furnishes the product R. The compound P upon treatment with an excess of phenol under basic conditions gives the product S.

The molar mass difference between compounds Q and R is 474 g mol⁻¹ and between compounds P and S is 172.5 g mol⁻¹.


[:Q.16] The number of heteroatoms present in one molecule of R is _____.

[Use: Molar mass (in g mol⁻¹): H = 1, C = 12, N = 14, O = 16, Br = 80, Cl = 35.5

Atoms other than C and H are considered as heteroatoms]

[:ANS] 9

[:SOLN]



Passage-2

A trinitro compound, 1,3,5-tris-(4-nitrophenyl)benzene, on complete reaction with an excess of Sn/HCl gives a major product, which on treatment with an excess of NaNO₂/HCl at 0 °C provides P as the product. P, upon treatment with excess of H₂O at room temperature, gives the product Q. Bromination of Q in aqueous medium furnishes the product R. The compound P upon treatment with an excess of phenol under basic conditions gives the product S.

The molar mass difference between compounds Q and R is 474 g mol⁻¹ and between compounds P and S is 172.5 g mol⁻¹.

[Q.17] The total number of carbon atoms and heteroatoms present in one molecule of S is _____.

[Use : Molar mass (in g mol⁻¹): H = 1, C = 12, N = 14, O = 16, Br = 80, Cl = 35.5

Atoms other than C and H are considered as heteroatoms]

[ANS] 51

[SOLN] From above diagram,

number of carbon atom in Structure S \cong Number of Phenylring \times 6 C-atom

\therefore total C-atom in Structure S = 7 \times 6 = 42

total number of hetero atom in S = 9